

## **Physics of the Earth results achieved at Department of Geophysics in 1999 (reported in May, 2000)**

A new fundamental monograph, summarizing outstanding research contributions of Prof. V. Červený to the ray theory of seismic waves has been completed (Červený, in press),

Development of the algorithms for two-point ray tracing and travel-time interpolation in 3D media has continued. The first quantitative results have been achieved in the description of the ray chaos due to heterogeneities in the velocity model by means of the average Lyapunov exponents. Various kinds of the coupling ray theory for weakly anisotropic models have been studied and compared with the exact solution derived for the "twisted crystal" model. Particular attention has been devoted to the resolution of seismic inversion techniques, model fitting, conversion and smoothing with the application of medium correlation functions and Sobolev scalar products (Bulant 1999a,b; Bulant and Klimeš, 1999a,b; Klimeš, 1999).

Theoretical modelling has closely touched seismic exploration problems. Ray and finite-difference methods have been used to simulate a measured CDP seismic section for the shallow lignite deposit at the Domenico lignite site, Greece (Brokešová et al., submitted). An important theoretical problem of inhomogeneous waves in dissipative media has been studied in detail. Extensive calculations of reflection/transmission coefficients of inhomogeneous waves at plane interfaces in various dissipative models have been performed and compared with the corresponding coefficients in elastic non-dissipative media (Brokešová, in press).

Theoretical computations of Fresnel volumes of PKP Earth's core waves have provided a suitable tool for estimating resolution power of tomographic method (Kvasnička and Janský, 1999). The distribution of body wave amplitudes and strains inside the Earth's mantle caused by an earthquake has been simulated (Duda et al., in press). Evaluation of the velocity models and relocation of weak earthquakes in the Western Bohemia earthquake swarms region have been performed by innovated methods (Janský, in press; Janský et al., in press; Málek et al., submitted).

The dispersion relation for Love waves in a layer on a half-space has been modified by introducing the wave number and its square instead of the phase velocity. The implicit function theorem has then been used to derive the analytical formulae for the group velocity and for the phase- and group-velocity partial derivatives with respect to the parameters of the medium. In comparison with the traditional formulation of the dispersion relation, this method is simpler and faster (Novotný, 1999).

New finite-difference methods for numerical simulation of seismic waves have been developed. The attention has been focused on stabilization procedures and the memory and time saving by means of irregular grids (Opršal and Zahradník, 1999a,b). A hybrid method has been developed and applied to earthquakes at EUROSEISTEST near Thessaloniki where it allowed a more realistic modeling than the conventional methods with plane waves (Riepl et al., 2000). The hybrid method has been supplemented by a stochastic component and used for the seismic-response analysis of the Tiber valley in Rome (Caserta et al., 1999). Disastrous Kobe earthquake of 1995 has been investigated within an international experiment (Opršal et al., 1999; Zahradník, 1999). New ASPO method for the focal-mechanism retrieval from earthquake amplitude spectra and polarities has been developed and applied in Corinth Gulf (Zahradník et al., in press). Seismic stations of the Charles University, operating in Greece, have contributed to the explanation of the damaging Athens 1999 earthquake (Tselentis and Zahradník, in press a,b).

Viscoelastic response of elastically compressible Earth's models has been studied. It has been found that unstable modes can exist (Hanyk et al., 1999). In the contrary, incompressible viscoelastic models have been found stable. The method of lines has been incorporated to discretize the system in space and a new eigenvalue problem for time evolution of viscoelastic models was obtained (Hanyk et al., submitted). This approach yields a complete spectrum of normal modes of viscoelastic relaxation without solving a secular equation in the Laplacian domain. The gravitational viscoelastic relaxation of eccentrically nested spheres has been solved semi-analytically in order to provide test examples for other numerical techniques (Martinec, 1999b; Martinec and Wolf, 1999).

Classical Boussinesq approximation of thermal convection was employed in simulations of thermal convection by Vecsey and Matyska (submitted). They studied wavelet spectra of temperature, kinetic energy and surface heat flow time series and clearly demonstrated the multiscale temporal dynamics of the convecting system. The extended-Boussinesq convection models have been used to extract the Bullen parameter. It has been shown that the profiles of the Bullen parameter have a definite potential of being useful in constraining the physical

parameters and flow structures associated with the Earth's mantle convection (Matyska and Yuen, 2000). Numerical simulations of the magnetohydrodynamic system have demonstrated that adiabatic heating/cooling stabilizes convection pattern in the Earth's core (Velínský and Matyska, 2000). The effects of adiabatic heating/cooling and viscous dissipation on 3-D rapidly rotating thermal convection was also studied by Mistr et al. (submitted).

The investigation of the Earth's mantle rheology has concentrated on forward and inverse analysis of the long-wavelength geoid in conjunction with seismic tomographic information and tectonic data (Čadek and Fleitout, 1999). It has been demonstrated that the available information can only be explained by the partially layered flow model that shows a strong increase of viscosity with depth and a pronounced asthenosphere below oceanic regions. At present, the preferred viscosity model is tested against the models of seismic anisotropy in the upper mantle. Thermal and mechanical coupling between the upper and the lower mantle was studied by Čížková et al. (1999), who showed that a low viscosity zone just beneath the 670 km boundary can effectively suppress the mechanical coupling and increase thus the importance of the thermal coupling.

The problem of the electromagnetic induction in a spherical Earth has been formulated in a weak sense and solved by the spectral-finite element technique (Martinec, 1999a). The spherical harmonic analysis has been carried out for magnetic storms within the range of periods of 5 to 40 days (Pruša and Martinec, 1999).

The theory of determination of precise geoid has been further developed in (Vaniček et al., 1999).

## References

- Brokešová, J., 2000. Reflection-transmission coefficients at a plane interface in dissipative and non-dissipative isotropic media: comparison. *J. Comp. Acoustics*, in press.
- Brokešová, J., Zahradník, J., and Paraskevopoulos, P. Ray and finite-difference modelling of CPD seismic sections for shallow lignite deposits. *J. Appl. Geophys.*, submitted.
- Bulant, P., 1999a. Two-point ray-tracing and controlled initial-value ray-tracing in 3-D heterogenous block structures. *Journal of Seismic Exploration*, **8**, 57-75.
- Bulant, P., 1999b. Calculation of multivalued ray-theory travel times in 3D – interpolation within individual ray cells. *Extended Abstracts of 61th EAGE Conference (Helsinki)*, Eur. Assoc. Geoscientists & Engr., Houten, P081.
- Bulant, P., Klimeš, L., 1999a. Interpolation of ray theory traveltimes within ray cells. *Geophysical Journal International*, **139**, 273-282.
- Bulant, P., Klimeš, L., 1999b. Interpolation within ray cells. *Expanded Abstracts of 69th Annual Meeting (Houston)*, Soc. Explor. Geophysicists, Tulsa, 1755-1758.
- Caserta, A., Zahradník, J., Plicka, V., 1999. Ground motion modelling with a stochastically perturbed excitation. *J. Seismology*, **3**, 45-59.
- Čadek, O., Fleitout, L., 1999. A global geoid model with imposed plate velocities and partial layering. *J. Geophys. Res.*, **104**, 29055-29076.
- Červený, V., 2000. Seismic Ray Theory. Cambridge Univ. Press, in press.
- Čížková, H., Čadek, O., van den Berg, A.P., Vlaar, N.J., 1999. Can lower mantle slab-like anomalies be explained by thermal coupling between the upper and lower mantles? *Geophys. Res. Lett.*, **26**, 1501-1504.
- Duda, S. J., Janský, J., Kvasnička, M. P wave amplitudes and dynamic strains inside the Earth. *Acta Geophysica Polonica*, in press.
- Hanyk, L., Matyska, C., Yuen, D.A., 1999. Secular gravitational instability of a compressible viscoelastic sphere. *Geophys. Res. Lett.*, **26**, 557-560.
- Hanyk, L., Matyska, C., Yuen, D.A. Eigenvalue approach in grid space to the problem of viscoelastic relaxation. *Geophys. Res. Lett.*, submitted.
- Janský, J. Grid search hypocentral location method in simple 1-D media. *Acta Montana*, in press.
- Janský, J., Horálek, J., Málek, J., Bousková, A. Homogeneous velocity models of the West Bohemian swarm region obtained by grid search. *Studia geoph. et geod.*, in press.
- Klimeš, L., 1999. Calculation of the third and higher travel – time derivatives in isotropic and anisotropic media. *Expanded Abstracts of 69th Annual Meeting (Houston)*, Soc. Explor. Geophysicists, Tulsa, 1751-1754.
- Kvasnička, M., Janský, J., 1999. Fresnel volumes corresponding to PKP waves in the IASP91 model. *Journal of Seismology*, **3**, 375-391.

- Málek, J., Horálek, J., Janský, J. Anisotropic velocity model of the Western Bohemia. *Phys. Earth Planet. Inter.*, submitted.
- Mistr, Z., Matyska, C., Yuen, D.A. A higher-order finite-difference method applied to rapidly rotating 3-D thermal convection at finite Prandtl number using the variables of vorticity, vector potential and temperature. *Geophysical Journal International*, submitted.
- Martinec, Z., 1999a. Spectral-finite element approach to three-dimensional electromagnetic induction in a spherical Earth. *Geophys. J. Int.*, **136**, 229-250.
- Martinec, Z., 1999b. Spectral, initial value approach for viscoelastic relaxation of a spherical Earth with a three-dimensional viscosity I. Theory. *Geophys. J. Int.*, **137**, 469-488.
- Martinec, Z., Wolf, D., 1999. Gravitational viscoelastic relaxation of eccentrically nested spheres. *Geophys. J. Int.*, **138**, 45-66.
- Matyska, C., Yuen, D.A., 2000. Profiles of the Bullen parameter from mantle convection modelling. *Earth Planet. Sci. Lett.*, **178**, 39-46.
- Novotný, O., 1999. A fast method of computing group-velocity partial derivatives for Love waves propagating in a layer on a half-space. *Studia geoph. et geod.*, **43**, 78-86.
- Opršal, I., Zahradník, J., 1999a. Elastic finite-difference method for irregular grids. *Geophysics*, **64**, 240-250.
- Opršal, I., Zahradník, J., 1999b. From unstable to stable seismic modelling by finite-difference method. *J. Phys. and Chem. of the Earth*, **24** (part A), no.3, 247-252.
- Opršal, I., Plicka, V., Zahradník, J., 1999. Kobe simulations by hybrid methods. In Irikura, K. et al. (eds.), *The effects of surface geology on seismic motion, Balkema, Rotterdam, Third volume, pp 1451-1456 (Proceedings of EGS 98, December 1-3 1998, Yokohama, Japan)*.
- Průša, L., Martinec, Z., 1999. Spherical harmonic analysis of geomagnetic variations observatory data up to degree 4 in the range of periods from 5 to 40 days. *Phys. Earth Planet. Inter.*, **115**, 229-245.
- Riepl, J., Zahradník, J., Plicka, V., and Bard, P. -Y., 2000. About efficiency of numerical 1-D and 2-D modelling of site effects in basin structures. *Pure and Appl. Geophys.*, **157**, 319-342.
- Tselentis, G.-A., Zahradník, J., 2000a. Aftershock monitoring of the Athens earthquake of September 7, 1999. *Seism. Res. Letters*, in press.
- Tselentis, G. -A., and Zahradník, J., 2000b. The Athens earthquake of September 7, 1999. *Bull. Seism. Soc. Am.*, in press.
- Vaniček, P., Huang, J., Novák, P., Pagiatakis, S., Véronneau, M., Martinec, Z., Featherstone, W.E., 1999. Determination of the boundary values for the Stokes-Helmert problem. *J. Geod.*, **73**, 180-192.
- Vecsey, L., Matyska, C. Wavelet spectra and chaos in thermal convection modelling. *Geophys. Res. Lett.*, submitted.
- Velínský, J., Matyska, C., 2000. The influence of adiabatic heating/cooling on magnetohydrodynamic systems. *Phys. Earth Planet. Inter.*, **117**, 197-207.
- Zahradník, J., 1999. Simulations of Kobe 1995 earthquake, reviewed in the light of previous and next international experiments. In Irikura, K. et al. (eds.), *The effects of surface geology on seismic motion, Balkema, Rotterdam, Third volume, pp 1503-1511 (Proceedings of EGS 98, December 1-3 1998, Yokohama, Japan)*.
- Zahradník, J., Hisada, Y., 1999. Nonlinearity and recent advancement in numerical simulations. In Irikura, K. et al. (eds.), *The effects of surface geology on seismic motion, Balkema, Rotterdam, Third volume, pp 1521-1522 (Proceedings of EGS 98, December 1-3 1998, Yokohama, Japan)*.
- Zahradník, J., Janský, J., and Papatsimpa, N., 2000. Focal mechanisms of weak earthquakes from amplitude spectra and polarities, *Pure and Appl. Geophys.*, in press.